

# $L_2$ distance between Gaussian Mixture Models

Kisung You

kisungyou@outlook.com

June 20, 2021

## 1 Introduction

Point set refers to a collection of points or observations. When we have multiple of such objects in a common ambient space, including  $\mathbb{R}^d$ , it is often convenient to represent each point set as a mixture of well-known probability distributions. This enables us to consider point sets as measures so that the analysis becomes operations on the space of measures.

In this note, we focus on a mixture of Gaussians, also known as Gaussian mixture model (GMM), and a measure of distance in the sense of  $L_2(\mathbb{R}^d)$  since it gives us explicit formula to compare two GMM-modeled measures  $P$  and  $Q$ . Existence of explicit expression makes it a fascinating option compared to other measures which may depend on Monte Carlo integration.

## 2 Problem Statement

Suppose we have two mixtures of Gaussians

$$P(x) = \sum_{n=1}^N \alpha_n N(x|\mu_n, \Sigma_n) \quad \text{and} \quad Q(x) = \sum_{m=1}^M \beta_m N(x|\eta_m, \Lambda_m)$$

where  $\alpha_n$  and  $\beta_m$  are nonnegative (actually, positive) coefficients that sum to 1 respectively, i.e.,  $\sum_{n=1}^N \alpha_n = 1$  and  $\sum_{m=1}^M \beta_m = 1$ .  $N(\cdot|\mu, \Sigma)$  and  $N(\cdot|\eta, \Lambda)$  are Gaussian distributions in  $\mathbb{R}^d$  with mean vectors  $\mu, \eta$  and covariance matrices  $\Sigma, \Lambda$  respectively. As everyone reading this note is probably well aware, the density function for  $N(x|\mu, \Sigma)$  is in form of

$$f(x) = \frac{1}{|\Sigma|^{1/2} (2\pi)^{d/2}} \exp\left(-\frac{1}{2}(x - \mu)^\top \Sigma^{-1}(x - \mu)\right)$$

with  $|\Sigma|$  being determinant of square covariance matrix  $\Sigma$ .

Two distributions are endowed with densities  $p(x)$  and  $q(x)$  for  $P$  and  $Q$  thanks to the above parametric forms and they are smooth functions over  $\mathbb{R}^d$ . This enables to define  $L_2$  distance between two distributions,

$$L_2(P, Q) = \left\{ \int_{\mathbb{R}^d} (p(x) - q(x))^2 dx \right\}^{1/2}. \quad (1)$$

For notational simplicity, we will denote  $p_n(x)$  and  $q_m(x)$  for densities of  $P_n$  and  $Q_m$  respectively.

### 3 Derivation

We want to compute  $L_2$  distance between two distributions  $P$  and  $Q$ . This can be written as

$$\begin{aligned}
 L_2^2(P, Q) &= \int_{\mathbb{R}} (p(x) - q(x))^2 dx \\
 &= \int \left( \sum_{n=1}^N \alpha_n p_n(x) - \sum_{m=1}^M \beta_m q_m(x) \right)^2 dx \\
 &= \sum_{n, n'} \alpha_n \alpha_{n'} \int p_n(x) p_{n'}(x) dx + \sum_{m, m'} \beta_m \beta_{m'} \int q_m(x) q_{m'}(x) dx \\
 &\quad - 2 \sum_{n, m} \alpha_n \beta_m \int p_n(x) q_m(x) dx
 \end{aligned}$$

which requires to evaluate integral for a product of two Gaussian densities,

$$\int p_n(x) q_m(x). \tag{*}$$

From the formula to be introduced in the next section, we have following equalities,

$$\begin{aligned}
 A_{n, n'} &= \int p_n(x) p_{n'}(x) dx = N(\mu_n | \mu_{n'}, \Sigma_n + \Sigma_{n'}) \\
 B_{m, m'} &= \int q_m(x) q_{m'}(x) dx = N(\eta_m | \eta_{m'}, \Lambda_m + \Lambda_{m'}) \\
 C_{n, m} &= \int p_n(x) q_m(x) dx = N(\mu_n | \eta_m, \Sigma_n + \Lambda_m).
 \end{aligned}$$

Therefore,

$$L_2(P, Q) = \left\{ \sum_{n, n'} \alpha_n \alpha_{n'} A_{n, n'} + \sum_{m, m'} \beta_m \beta_{m'} B_{m, m'} - 2 \sum_{n, m} \alpha_n \beta_m C_{n, m} \right\}^{1/2}$$

#### Fact (\*)

According to Section 8.1.8 of Matrix Cookbook [1], product of two gaussian densities is in form of following,

$$N(x | m_1, \Sigma_1) \cdot N(x | m_2, \Sigma_2) = c_c \cdot N(x | m_c, \Sigma_c)$$

where

$$\begin{aligned}
 c_c &= N(m_1 | m_2, (\Sigma_1 + \Sigma_2)) \\
 m_c &= (\Sigma_1^{-1} + \Sigma_2^{-1})^{-1} (\Sigma_1^{-1} m_1 + \Sigma_2^{-1} m_2) \\
 \Sigma_c &= (\Sigma_1^{-1} + \Sigma_2^{-1})^{-1}.
 \end{aligned}$$

From the fact that integral of density function is 1, we have

$$\int N(x | m_1, \Sigma_1) \cdot N(x | m_2, \Sigma_2) dx = c_c.$$

### References

- [1] K. B. Petersen and M. S. Pedersen. The matrix cookbook, nov 2012. Version 20121115.